

# A new species of *Ischiodon* Sack (Diptera, Syrphidae) from Madagascar

Ximo Mengual<sup>1</sup>

<sup>1</sup> Zoologisches Forschungsmuseum Alexander Koenig, Leibniz-Institut für Biodiversität der Tiere, Adenauerallee 160, D-53113 Bonn, Germany

Corresponding author: Ximo Mengual ([x.mengual@leibniz-zfmk.de](mailto:x.mengual@leibniz-zfmk.de))

---

Academic editor: B. Müller | Received 14 February 2018 | Accepted 9 April 2018 | Published 18 April 2018

---

<http://zoobank.org/BCA6F985-BE53-47C5-B64D-005DA0DD69BE>

---

**Citation:** Mengual X (2018) A new species of *Ischiodon* Sack (Diptera, Syrphidae) from Madagascar. African Invertebrates 59(1): 55–73. <https://doi.org/10.3897/AfrInvertebr.59.24461>

---

## Abstract

The flower fly genus *Ischiodon* Sack (Diptera, Syrphidae) is revised and a new species, *Ischiodon astales* sp. n., is described from Madagascar. Additionally, a lectotype for *Ischiodon aegyptius* is designated and the first records of *Ischiodon scutellaris* for the Arabian Peninsula are reported. Diagnoses, illustrations, synonymies and distributional data are given for all described species, as well as an identification key to all known species.

## Keywords

identification key, flower flies, hoverflies, *Ischiodon astales*, Afrotropical Region

## Introduction

The genus *Ischiodon* Sack, 1913 (Diptera, Syrphidae) is a small, but very characteristic genus, with a very broad distribution and a taxonomic history full of confusion and synonyms (Vockeroth 1969). The genus occurs in southern Europe, the Mediterranean basin, throughout all of the African continent, eastwards to India, China, Japan, southeast Asia, Australia and some Pacific islands (Huang and Cheng 2012, Thompson 2013, Speight 2017). *Ischiodon* species have been originally described in eight different genera, a fact that may indicate the morphological resemblance of these

species with other syrphid taxa and the broad genus concepts of previous authors: i.e. *Musca* Linnaeus, 1758, *Syrphus* Fabricius, 1775, *Scaeava* Fabricius, 1805, *Sphaerophoria* Le Peletier & Serville, 1828, *Xanthogramma* Schiner, 1860, *Epistrophe* Walker, 1852, *Melithreptus* Loew, 1840 (a junior primary homonym of *Sphaerophoria*), and *Ischiodon* (Thompson 2013). On the other hand, a junior synonym of *Sphaerophoria rueppellii* (Wiedemann, 1830) was described as *Ischiodon libicum* Nayar, 1978.

The confusion of *Ischiodon* as a valid genus is due to its superficial morphological resemblance with the genus *Simosyrphus* Bigot, 1882. This confusion started when Bezz (1912:72) synonymised *Syrphus grandicornis* Macquart, 1842 (now *Simosyrphus grandicornis*) under *Ischiodon scutellaris* (Fabricius, 1805). Species of both genera have been previously placed under one another several times (Mengual 2015, see below under each species), despite some important differences in the male genitalia (Vockeroth 1969). The generic key by Vockeroth (1969) can be used to identify both genera. A synonymy of both genera was proposed by Láska et al. (2006) based on larval and pupal morphology, supported by a molecular study of a very limited taxon sampling using only part of the gene *cytochrome c oxidase subunit 1* (COI). More recently, Mengual et al. (2018) resolved both genera as related to *Scaeava* but the genus *Pseudodoros* as sister group of *Ischiodon*; thus, current evidence supports the generic rank of *Ischiodon* and *Simosyrphus*.

*Ischiodon* comprises three species with quite distinct distributions: *I. feae* (Bezz, 1912) is endemic of the Cape Verde Islands, *I. aegyptius* (Wiedemann, 1830) is mainly present in Africa and southern Europe and *I. scutellaris* (Fabricius, 1805) occurs in the Oriental and Australasian Regions. Larvae of *I. aegyptius* and *I. scutellaris* have been reported feeding on aphids (Hemiptera, Aphididae) mostly, but there are a few records on thrips (Thysanoptera), whiteflies (Hemiptera, Aleyrodidae) and caterpillars (Lepidoptera) (Rojo et al. 2003).

The aim of this study is to revise the genus *Ischiodon* and describe a new species from Madagascar, as well as to provide an identification key for all known species of this genus. In addition, a lectotype is designated for *Syrphus aegyptius* Wiedemann, 1830 and the first records of *I. scutellaris* from the UAE are reported.

## Material and methods

Differential diagnoses, synonymies and distributions are given for all studied species. The new species is described in full, with terminology following Thompson (1999) and Cumming and Wood (2009). The abbreviations used for collections follow the standard of the *Systema Dipterorum* (Thompson 2013) and their equivalents are given below:

<b>CAS</b>	California Academy of Sciences, San Francisco, USA.
<b>CSCA</b>	California State Collection of Arthropods, Sacramento, USA.
<b>MSNG</b>	Museo Civico di Storia Naturale ‘Giacomo Doria’, Genova, Italy.
<b>MZH</b>	Finnish Museum of Natural History, Helsinki, Finland.

- SMF** Senckenberg Forschungsinstitut und Naturmuseum, Frankfurt am Main, Germany.  
**ZFMK** Zoologisches Forschungsmuseum Alexander Koenig, Bonn, Germany.  
**ZMUC** University of Copenhagen, Zoological Museum, Copenhagen, Denmark.

In the description of type labels, the contents of each label are enclosed within double quotation marks (" "), italics denote handwriting and the individual lines of data are separated by a double forward slash (//). At the end of each record, between square brackets ([ ]) and separated by a comma, the number of specimens and sex, the holding institution and the unique identifier or number are given.

All measurements are in millimetres and were taken using a reticule in a Leica M165 C microscope. Photographs were composed using the software Zerene Stacker 1.04 (Richland, Washington, USA), based on images of pinned specimens taken with a Canon EOS 7D mounted on a P-51 Cam-Lift (Dun Inc., VA, USA) and with the help of Adobe Lightroom (version 5.6). Simple-Mappr (Shorthouse 2010) was used to create Figure 4. Body length was measured from the anterior oral margin to the posterior end of the abdomen, in lateral view. Wing length was measured from the wing tip to the basicosta.

## Results

### Genus *Ischiodon* Sack, 1913

**Type-species.** *Ischiodon trochanterica* Sack, 1913, by monotypy; junior synonym of *Scaeva scutellaris* Fabricius, 1805.

**Differential diagnosis** (adapted from Vockeroth 1969). Small to medium-sized, rather slender species with yellow-margined mesonotum, broad yellow fasciae on abdomen and very large and prominent male genitalia. Face yellow, with or without medial black vitta; eye bare; basoflagellomere large, twice as long as broad, tapering slightly to an acutely rounded apex. Mesonotum shining black with broad, well-defined, pale to bright yellow lateral vitta extending to postalar callus. Scutellum yellow, usually obscurely brownish on disc. Pleuron mostly shining black, except posterior anepisternum yellow on posterior half and katepisternum with a dorsal yellow macula. Dorsal and ventral katepisternal pile patches broadly separated posteriorly. Metasternum bare. Wing with microtrichia greatly reduced, absent from most of basal half and elsewhere very sparse and irregularly scattered. Abdomen elongate, parallel-sided or very narrowly oval, rather flattened, with a strong margin from middle of tergite 2 to the end of tergite 5. Tergite 2 with a pair of separate or narrowly confluent pale maculae; tergites 3 and 4 each with a rather broad, slightly arcuate yellow fascia separated from the base of each tergite. Very often, the black abdominal areas become reddish or yellow towards the abdominal apex, so the pattern is indistinct.

In males of *I. aegyptius* and *I. scutellaris*, the ventral surface of the metatrochanter has a slender or moderately stout, cylindrical, apically acute process (calcar) of varying length. In males of *I. aegyptius* and *I. feae*, the inner (mesial) claw of the protarsus has a dorsal preapical projection, or flange, which gives the claw a truncated appearance (Fig. 5F).

**Etymology.** *Ischiodon* is derived from the neuter adjective *ischion*, from the Greek *ischion* meaning “hip, coxa” (Brown 1956: 411) and the masculine Greek noun *odous* (*odon*), *odontos*, meaning tooth (Brown 1956: 807). Thus, *Ischiodon* refers to the calcar or spur that some species have in the metatrochanter.

Several authors (including the present author) used the species epithet of *Ischiodon aegyptium* in neuter form. *Ischiodon* must be treated as a masculine gender and all species names must follow this gender except nouns in apposition. Thus, the correct species epithet is *Ischiodon aegyptius*.

### *Ischiodon aegyptius* (Wiedemann, 1830)

Figure 1

*Musca nigra* Forsskål, 1775: xxiv. Type-locality: Egypt, Arabia. Type material most likely lost (not found in the digitised type collection of the ZMUC).

*Syrphus aegyptius* Wiedemann, 1830: 133. Type-locality: Eritrea and Ethiopia, as Abyssinia, here restricted. Lectotype ♂, SMF, here designated.

*Syrphus senegalensis* Guérin-Méneville, 1832: pl. 99. [Description in 1844: 545]. Type-locality: Senegal.

*Sphaerophoria annulipes* Macquart, 1842: 103. Type-locality: Egypt.

*Syrphus longicornis* Macquart, 1842: 154. Type-locality: South Africa.

*Syrphus natalensis* Macquart, 1846: 262 (134). Type-locality: South Africa, ‘Port Natal’.

*Syrphus felix* Walker, 1852: 229. Type-locality: Canary Is.


*Syrphus brachypterus* Thomson, 1869: 496. Type-locality: Madeira.

*Sphaerophoria pyrrura* Bigot, 1884: 99. Type-locality: Senegal.

*Sphaerophoria borbonica* Bigot, 1884: 100. Type-locality: Réunion Is.

*Xanthogramma catalonicum* Andreu, 1926: 110. Type-locality: Spain: Prov. Barcelona, Moya.

**Differential diagnosis.** Very common species in the Afrotropical region with yellow markings on abdomen reaching the lateral margins (Fig. 1A, C). Males of *I. aegyptius* have metatrochanters with a long calcar (Fig. 1E) (short calcar in *I. scutellaris*) and the claws of the fore leg are asymmetrical (as in Fig. 5F) (males of *I. feae* also have asymmetrical claws but they do not have a calcar on the metatrochanters). The male genitalia are quite unique with a trilobed surstyli (Fig. 1F). Females are very difficult to distinguish from females of *I. scutellaris* as the pattern of tergite 2, very often used to distinguish them (Sack 1932, Huang and Cheng 2012, Speight and Sarthou 2017), is variable. At this moment, conspecific males, DNA barcodes and/or collecting locality may help to distinguish them.



**Figure 1.** *Ischiodon aegyptius* (Wiedemann), male ZFMK-DIP-00012424: **A** Habitus, lateral view **B** Frontal view **C** Habitus, dorsal view **D** Labels **E** Metacoxa, metatrochanter and metafemur, lateral view **F** Male genitalia, lateral view (from Claussen and Barkemeyer 1987). Scale bars: 1 mm.

**Geographical distribution.** African continent and archipelagos, southern Europe, Israel and Arabian Peninsula.

**Type-locality.** Wiedemann (1830) studied an unknown number of specimens from Egypt and Nubia (currently divided into Egypt and Sudan). All the syntypes present in the ZMF collection, collected by Rüppell with original handwritten labels, were collected from Abyssinia, a historical region in northern Ethiopia and southern Eritrea, although sometimes east Sudan is also considered part of this kingdom. Thus, the type-locality is here restricted to Eritrea and Ethiopia based on the lectotype designation.

**Material examined.** More than 300 specimens from UAE, Israel, Cyprus, Morocco, Canary Islands, Madagascar, Kenya, The Gambia, Central African Republic, Zimbabwe and South Africa.

**Remarks.** Amongst the syntypes, deposited at the Senckenberg Forschungsinstitut und Naturmuseum (SMF, Frankfurt am Main, Germany), there is a pinned male labelled “Abyssinia // Dr. Rüppell.” “289” [green] “*Ischiodon* // *aegyptius* Wd” “Lecto- // Typus” [red] “Dipt. // 300” [reverse] “LECTOTYPE // *Ischiodon* // *aegyptius* // det. X. Mengual 2017” [red, second and third lines handwritten]. This specimen is here designated lectotype to fix and ensure the universal and consistent interpretation of the name. The other syntypes, a male and a female without head, have been labelled as paralectotypes: “Abyssinia // Dr. Rüppell.” “*Ischiodon* // *aegyptius* Wd” “Para- // typoid” [red] “Dipt. // 300” [reverse] “PARALECTOTYPE // *Ischiodon* // *aegyptius* // det. X. Mengual 2017” [yellow].

Some authors (Huang et al. 1996, Huang and Cheng 2012) reported specimens identified as *I. aegyptius* from China, with a complete overlapping distribution inside this country with *I. scutellaris* (see Huang and Cheng 2012). In the author’s opinion, only *I. scutellaris* occurs in China and these specimens are misidentifications when the pattern of the tergite 2 is used in females for identification. These works (Huang et al. 1996, Huang and Cheng 2012) do not specify if there are males of *I. aegyptius* amongst the studied material, which could be used to corroborate the author’s opinion.


### *Ischiodon astales* sp. n.

<http://zoobank.org/B32ED255-02F5-4465-8CBC-7B7EA64404CC>

Figures 2, 3A–D, 4

**Differential diagnosis.** Females and males of this new species do not have a calcar on the metatrochanter and they are easy to distinguish from other species since the two broad, yellow maculae on tergite 2 do not reach the lateral margins (Figs 2A, C, 3A, C). Moreover, males of *I. astales* sp. n. have symmetrical claws on the fore leg, the same as males of *I. scutellaris*.

**Description. Male:** Head (Fig. 2A, B): Face with distinct facial tubercle, entirely yellow, yellow pilose; gena yellow with two small black macula at eye margin; lunule yellow; frons yellow, yellow pilose; vertical triangle black; antenna yellow, black pilose, basoflagellomere orange, darker dorsally; arista orange-yellow, bare; eye bare; occiput



**Figure 2.** *Ischiodon astales* sp. n., male holotype CASENT 2210258: **A** Habitus, lateral view **B** Frontal view **C** Habitus, dorsal view **D** Labels **E** Male genitalia, lateral view **F** Hypandrium, ventral view. Scale bars: 1 mm.

black, white pollinose and white pilose on ventral 1/2, yellow pollinose and yellow pilose on dorsal 1/2.

**Thorax** (Fig. 2A, C): Scutum shiny black, with long yellow pile; postpronotum paler, brownish, bare; notopleuron and supra-alar area yellow; postalar callus darker, brownish; scutellum yellow with medial area brownish, yellow pilose, subscutellar fringe yellow. Pleuron black, except posterior anepisternum yellow on posterior 1/2 and katepisternum with a dorsal yellow macula, yellow pilose; katepisternal pile patches separated; metaepisternum bare; metasternum bare; postmetacoxal bridge incomplete; calypter yellow, yellow pilose; plumula long, pale; halter pedicel and capitulum yellow; posterior spiracular fringes yellow. **Wing**: hyaline, bare on basal 1/2 and with sparse and irregularly scattered microtrichia on apical 1/2; alula mostly bare, with a few microtrichia posteriorly. **Legs**: fore and mid legs yellow with femora slightly more orange, coxae black and basal 1/2 of trochanters dark, entirely yellow pilose; hind leg with black coxae and basal part of trochanter dark; yellow metafemur with an almost imperceptible brown area on the apical 1/2, and metatibia slightly darker (dark yellow) on apical 1/2, black and yellow pilose.

**Abdomen** (Fig. 2A, C): Parallel-sided, with lateral margin from apical 1/2 of tergite 2 to tergite 5. Tergite 1 black, except yellow on lateral margins, yellow pilose; tergite 2 black, paler medially on posterior margin, with two large, rounded yellow maculae that do not reach lateral margins, yellow pilose, with short black pile on posterior margin; tergite 3 black on basal 1/2 becoming orange or brownish on posterior 1/2, with a medial yellow fascia that does not reach the lateral margins, with scattered, short, black pile; tergite 4 brownish-orange with a medial yellow fascia, black pilose except on the yellow fascia; tergite 5 brownish-orange, black pilose. Sternite 1 yellow; sternite 2 yellow with a medial, round, black macula; sterna 3 and 4 yellow becoming darker (brownish orange) towards genitalia; all sterna with yellow pile. Male genitalia: large, brownish-orange, as in Fig. 2E, F.

**Female**: Similar to male except for normal sexual dimorphism and as follows: frons black on dorsal 1/3, yellow on ventral 2/3 with a variable medial brownish orange vitta.

**Variation**: Due to the small number of specimens, one must expect variability in the coloration of the abdomen, legs and female frons, as in other *Ischiodon* species.


**Length** (N=4): Body, 10.0 mm (9.0–12.0 mm); wing, 7.7 mm (7.5–8.0 mm).

**Geographical distribution.** Species known from the southern regions of Atsimo-Andrefana and Anosy, part of the former Toliara Province, Madagascar (Fig. 4; see Suppl. material 1: Table S1).


**Etymology.** The specific epithet is derived from the Greek *astales* meaning unarmed, unclad (Brown 1956: 122). Species epithet is to be treated as an adjective.

**Type locality.** Madagascar: Atsimo-Andrefana Region, Mikea forest, NW of Manombo, 22°54.22'S, 43°28.53'E.

**Material examined.** Type material. *Holotype*, male, pinned with genitalia in microvial, deposited in the CAS and labelled: “MADAGASCAR, Tulear // Province, Mikea Forest, // NW of Manombo, el 30 m, // 17 – 27 July 2003” “22°54.22'S, 43°28.53'E // coll: M. Irwin, R. Harin’Hala // California Acad of Sciences // malaise trap - in deciduous //



**Figure 3.** *Ischiodon astales* sp. n., female paratype CASENT 2210121: **A** Habitus, lateral view **B** Frontal view **C** Habitus, dorsal view **D** Labels. *Ischiodon feae* (Bezzi), female GJ.1985: **E** Habitus, dorsal view **F** Frontal view. Scale bars: 1 mm (**A–C, E**); 0.5 mm (**F**).



**Figure 4.** Known distribution of *Ischiodon astales* sp. n.

dry forest MA-02-18A-62” “CASENT // 2210258” “HOLOTYPE // *Ischiodon* // *astales* // Det. X. Mengual 2017” [red]. Paratypes: Madagascar: Atsimo-Andrefana Region, Mikea Forest, NW of Manombo, 37 m., -22.91333°, 43.48217°, 12–20.xi.2001, M. Irwin, R. Harin’Hala [1♀, ZFMK; CASENT 2210121]; Atsimo-Andrefana Region, Zombitse-Vohibasia National Park, near ANGAP Office, 840 m., -22.8865°, 44.69217°, 31.x–16.xi.2002, R. Harin’Hala [1♀, CSCA; ZFMK-DIP-00026782]; Anosy Region, Berenty Special Reserve, 8 km NW Amboasary, 85 m., -25.00667°, 46.30333°, 26.x–2.xi.2002, M. Irwin, F. Parker, R. Harin’Hala [1♀, ZFMK; CASENT 2210162]; Anosy Region, Andohahela National Park, Ihazofotsy Parcelle III, 80 m., -24.83083°, 46.53617°, 23–30.xi.2003, M. Irwin, F. Parker, R. Harin’Hala [1♀, CAS; CASENT 2210150].

**Remarks.** All specimens of this new species were collected using Malaise traps in different habitats. The male holotype was collected in a deciduous dry forest, while females CASENT 2210150 and CASENT 2210121 were collected in dry spiny forests and the female CASENT 2210162 in a gallery forest. On the other hand, the female ZFMK-DIP-00026782 was collected at a relative high altitude compared with the other specimens.


### *Ischiodon feae* (Bezzi, 1912)

Figures 3E, F, 5

*Syrphus feae* Bezzi, 1912: 407. Type, MSNG. Type-locality: Cape Verde Is.

**Differential diagnosis.** Very distinct *Ischiodon* species with a darker overall coloration (Figs 3E, 5C), face yellow with a medial black vitta (Figs 3F, 5B) and terga 2–4 with two small, oblique yellow maculae not reaching the lateral margins (Fig. 5A, C). Males of *I. feae* have fore legs with asymmetrical claws and metatrochanter without calcar.

**Geographical distribution.** Only known from Cape Verde.



**Figure 5.** *Ischiodon feae* (Bezzi), male <http://id.luomus.fi/GJ.1984>: **A** Habitus, lateral view **B** Frontal view **C** Habitus, dorsal view **D** Labels **E** Wing **F** Acropod and fifth tarsomere of the fore leg, dorsal view **G** Male genitalia, lateral view **H** Hypandrium, ventral view. (**E-H** from Claussen and Barkemeyer 1987). Scale bars: 1 mm.

**Material examined.** CAPE VERDE: São Vicente Is., Monte Verde, 25–29.xi.1953, Lindberg [1♂, MZH; <http://id.luomus.fi/GJ.1984>]; São Vicente Is., Ribeira de Julião, 26.xi–2.xii.1953, Lindberg [1♀, MZH; <http://id.luomus.fi/GJ.1985>]; Santa Luzia Is., Água Doce, 3.xii.1953, Lindberg [2♀, MZH; <http://id.luomus.fi/GJ.1986>, <http://id.luomus.fi/GJ.1987>].

### *Ischiodon scutellaris* (Fabricius, 1805)

Figure 6

*Scaeva scutellaris* Fabricius, 1805: 252. Type-locality: India: Tamil Nadu, Tharangambadi. Syntypes, ZMUC.

*Syrphus coromandensis* Macquart, 1842: 149. Type-locality: India: ‘Cote de Coromandel’.

*Sphaerophoria annulipes* Macquart, 1855: 116. Type-locality: Marquesas Is.

*Syrphus splendens* Doleschall, 1856: 410. Type-locality: Indonesia: Java.

*Syrphus nodalis* Thomson, 1869: 497. Type-locality: French Polynesia: Society Is., Tahiti.

*Syrphus erythropygus* Bigot, 1884: 87. Type-locality: ‘Indes’.

*Syrphus ruficauda* Bigot, 1884: 96. Type-locality: New Caledonia.

*Melithreptus novaeguineae* Kertesz, 1899: 178. Type-locality: Papua New Guinea: Astrolabe Bay, Erima.

*Ischiodon trochanterica* Sack, 1913: 6. Type-locality: Taiwan: Kanshizei, Puli as ‘Polisha’, Suihenkyaku, Tainan and Takao.

*Melithreptus ogasawarensis* Matsumura, 1916: 23. Type-locality: Japan: Bonin Is., Ogasawara-jima.

*Ischiodon boninensis* Matsumura, 1919: 128. Type-locality: Japan: Bonin Is., Chichijima, Ogasawara-jima.

*Epistrophe platychiroides* Frey, 1946: 164. Type-locality: Philippines: Luzon, Quezon, Atimonan.


*Ischiodon penicillatus* Hardy, 1952: 363; *nomen nudum*.

*Epistrophe magnicornis* Shiraki, 1963: 141. Type-locality: Micronesia: Chuuk State, Weno Is.

*Sphaerophoria macquarti* Goot, 1964: 220. New name for *annulipes* Macquart.

**Differential diagnosis.** Very widespread species with yellow abdominal marking reaching lateral margins (Fig. 6A, C). Males have the metatrochanters with a short calcar (Fig. 6E) and the claws in all legs are symmetrical. The pattern of tergite 2 in females is variable and cannot always be relied on to distinguish the females of *I. scutellaris* from those of *I. aegyptius*.

**Geographical distribution.** From Greece, eastwards to Caucasus, Kazakhstan, Iran, Arabian Peninsula south to Indomalayan region, China, Japan, Taiwan and Australasian and Oceanian regions except Hawaii. *I. scutellaris* has not been reported from Korea (Han and Choi 2001).



**Figure 6.** *Ischiodon scutellaris* (Fabricius), male ZFMK-DIP-00019729: **A** Habitus, lateral view **B** Frontal view **C** Habitus, dorsal view **D** Labels **E** Metacoxa and metatrochanter, posterolateral view **F** Male genitalia: **a** lateral view **b** postgonite, outline only (from Vockeroth 1969). Scale bars: 1 mm.

**Material examined.** More than 150 specimens from Greece, UAE, Kazakhstan, Iran, Nepal, Indonesia, Thailand, Taiwan, Australia, New Caledonia and French Polynesia. The syntypes of *Scaeva scutellaris* Fabricius, 1805 were not studied for the present study.

**Remarks.** In their excellent treatment of the flower fly fauna of the Arabian Peninsula, Smit et al. (2017) recorded *I. aegyptius* (as *Simosyrphus aegyptius*) from the UAE, Yemen and Socrota Is. and mentioned that the species was previously known from Saudi Arabia. Amongst the examined material, there are two males of *I. scutellaris* labelled as follows: UAE: Wadi Bih, dam, 25°48'N, 56°4.20'E, 4–13.i.2011, C. Schmid-Egger [1♂, CSCA; ZFMK-DIP-00026784]; Wadi Maidaq, 25°18.6'N, 56°7.8'E, 11–19. iii.2009, C. Schmid-Egger [1♂, CSCA; ZFMK-DIP-00026785]. These two males are the first records of *I. scutellaris* for the UAE and for the entire Arabian Peninsula.

### Key to species of *Ischiodon* Sack, 1913

- |   |                                                                                                                                                                                                                                                                                                                                                                                                                                                               |                                |
|---|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------|
| 1 | Males: eyes meeting on frons (Figs 1B, 2B, 5B, 6B) .....                                                                                                                                                                                                                                                                                                                                                                                                      | 2                              |
| — | Females: eyes separated (Fig. 3B, F) .....                                                                                                                                                                                                                                                                                                                                                                                                                    | 5                              |
| 2 | Metatrochanter without a calcar .....                                                                                                                                                                                                                                                                                                                                                                                                                         | 3                              |
| — | Metatrochanter with a calcar (spur) (Figs 1E, 6E) .....                                                                                                                                                                                                                                                                                                                                                                                                       | 4                              |
| 3 | Face entirely yellow (Fig. 2B). Tergite 2 with two broad, rounded yellow maculae; terga 3 and 4 mostly orange (Fig. 2A, C). Claws on all legs normal, symmetrical. Male genitalia: surstylus broadened beyond base with convex, dorsal margin; postgonite dorsally with two tips oriented anteriorly (Fig. 2E, F) (Madagascar) .....                                                                                                                          | <i>astales</i> sp. n.          |
| — | Face yellow with a medial black vitta (Fig. 5B). Terga 2–4 with small, oblique yellow maculae (Fig. 5A, C). Claws of fore leg asymmetrical: inner claw with a rounded flange dorsally, just before tip, in addition to pointed tip (Fig. 5F). Male genitalia: surstylus elongated, parallel-sided; postgonite dorsally with two lobes, the anterior (basal) lobe with two tips (Fig. 5G, H) (Cape Verde) .....                                                | <i>feae</i> (Bezzi)            |
| 4 | Calcar on metatrochanter long, more than three times longer than wide (Fig. 1E). Claws of fore leg asymmetrical: inner claw with a rounded flange or projection dorsally, just before tip, in addition to pointed tip (Fig. 5F). Male genitalia: surstylus trilobed, with dorsal margin convex; postgonite with a dorsal, very long, slightly curved process (Fig. 1F) (African continent and archipelagos, southern Europe, Israel, Arabian Peninsula) ..... | <i>aegyptius</i> (Wiedemann)   |
| — | Calcar on metatrochanter shorter, about two times longer than wide (Fig. 6E). Claws on all legs normal, symmetrical. Male genitalia: surstylus broadened beyond base with convex, weakly serrate, dorsal margin; postgonite elongated with a broad base and a small process (Fig. 6F) (from Greece, Caucasus, Kazakhstan, Iran, Arabian Peninsula south to Indomalayan Region, Australasian and Oceanian regions, Japan).....                                 | <i>scutellaris</i> (Fabricius) |

- 5 Yellow abdominal maculae on terga 2 and 3 not reaching lateral margins (Fig. 3A, C, E) ..... 6
- Yellow abdominal maculae on terga 2 and 3 reaching lateral margins ..... 7
- 6 Face entirely yellow (Fig. 3B). Abdomen dark anteriorly, but terga 3–6 reddish; tergite 2 with two large yellow maculae and terga 3 and 4 each with a yellow fascia (Fig. 3A, C) (Madagascar) ..... *astales* sp. n.
- Face yellow with a medial black vitta (Fig. 3F). Abdomen dark; terga 2–4 each with two oblique yellow maculae (Fig. 3E) (Cape Verde) ..... *feae* (Bezzi)
- 7 Typical female: tergite 2 usually with a transverse yellow fascia, sometimes with two yellow maculae with pointed inner margin (African continent and archipelagos, southern Europe, Israel, Arabian Peninsula) ..... *aegyptius* (Wiedemann)
- Typical female: tergite 2 usually with two yellow maculae with rounded inner margin (from Greece, Caucasus, Kazakhstan, Iran, Arabian Peninsula south to Indomalayan Region, Australasian and Oceanian regions, Japan) ..... *scutellaris* (Fabricius)
- NOTE: Females of *I. aegyptius* and *I. scutellaris* cannot be distinguished as the pattern of tergite 2 is very variable. At this moment, conspecific males, DNA barcodes and/or collecting locality may help to distinguish them.

## Discussion

A new species of *Ischiodon* is described from the southern regions of Madagascar. It should be noted that the male holotype and the four female paratypes of *I. astales* sp. n. were found amongst more than 200 specimens of *Ischiodon aegyptius* collected over several years (2002–2008) using Malaise traps (specimens deposited in the CAS). This might indicate the scarcity of this new species in nature, but the fact that *I. astales* has been collected in several habitats and at different elevations might indicate the opposite, i.e. the new species is more common than the number of specimens studied suggests.

*Ischiodon scutellaris* has reached the Mediterranean Basin and it is present in Greece, Crimea Peninsula and Turkey and might eventually reach northern areas of the Balkan Peninsula. In contrast, the species present in the Iberian Peninsula, Macaronesia, Italy, northern parts of the African continent, Cyprus and Israel is *I. aegyptius*. It seems that there is an overlapping area of these two species including the southern region of Turkey and Iran, where both species have been reported (Kazerani et al. 2013, Naderloo et al. 2013, Saribiyik 2014, Khosravian et al. 2015) and now also in the UAE. One is biased to think that the specimens of *I. scutellaris* collected in the UAE arrived from across the Persian Gulf or the Gulf of Oman, as there are no records of *I. scutellaris* from Iraq, Kuwait or Saudi Arabia. On the other hand, *I. scutellaris* might be present in these countries, but the lack of data makes any assumption difficult to corroborate as the syrphid fauna from these countries is poorly known.

Speight (2017) affirms that *I. aegyptius* is possibly a non-resident species in Europe, but it repeatedly establishes temporary populations that may, or may not, survive from one year to the next in Europe. Contrary to Speight (2017), the author believes that *I. aegyptius* and *I. scutellaris* are both resident species in Europe based on the number of continuous records and new evidence, as the habitats in the Mediterranean Basin between countries are not so different.

## Acknowledgements

My most sincere thanks to Martin Hauser for sharing with me the information about syrphid fauna from the UAE. Thanks to Gunilla Ståhls (MZB), Martin Hauser (CSCA), Irene Rademacher and Andrea Hostenpflug-Vesmanis (SMF) and Michelle Trautwein (CAS) for letting me study the material in their care. I am grateful to Claus Claussen, Eyjolf Aistleitner and Werner Barkemeyer for advice and help regarding *I. feae* material and for letting me use their drawings and to Christodoulos Makris for his help in the field in Cyprus and for the advice on the Greek language. I thank John T. Smit, Gunilla Ståhls and Gil Felipe G. Miranda for their very helpful comments. I also thank Trevor O. Burt for the excellent drawing of male genitalia of the new species *I. astales*.

## References

- Andreu J (1926) Notas dipterológicas. Una lista de Sírfidos para contribuir al conocimiento de los dípteros de España. Boletín de la Sociedad Entomológica de España 9: 98–126.
- Bezzi M (1912) Ditteri raccolti da Leonardo Fea durante il suo viaggio nell' Africa occidentale. Parte 1a: Syrphidae. Annali del Museo Civico di Storia Naturale di Genova 45: 400–453.  
<https://biodiversitylibrary.org/page/10812452>
- Bigot JMF (1882) Annales de la Société Entomologique de France 2(6): 67–68. <https://biodiversitylibrary.org/page/8997996> [Descriptions de quatre genres nouveaux de la tribu des Syphides (Syrphidae auctorum), ainsi que celles de deux nouvelles espèces]
- Bigot JMF (1884) Dipteres nouveaux ou peu connus. 24 partie, XXXII: Syrphidi (2 partie). Espèces nouvelles, no. III. Annales de la Société entomologique de France ser 6(4): 73–80, 81–116. <https://biodiversitylibrary.org/page/32550189>
- Brown RW (1956) Composition of scientific words a manual of methods and a lexicon of materials for the practice of logotechnics. Published by the author, Baltimore, 882 pp. <https://archive.org/details/compositionofsci00brow>
- Claussen C, Barkemeyer W (1987) Zur Syrphidenfauna der Kapverden (Insecta: Diptera: Syrphidae). Courier Forschungsinstitut Senckenberg 95: 71–86.
- Cumming JM, Wood DM (2009) Adult morphology and terminology. In: Brown B, Bor-kent A, Cumming JM, Wood DM, Woodley NE, Zumbado MA (Eds) Manual of Central American Diptera, Vol. 1. NRC Press, Ottawa, 9–50.

- Doleschall CL (1856) Eerste bijdrage tot dekennis der dipterologische fauna van Nederlandsch Indie. Natuurkundig Tijdschrift voor Nederlandsch Indië 10: 403–414. [12 pls] <https://biodiversitylibrary.org/page/13612961>
- Fabricius JC (1775) Systema entomologiae: sistens insectorum classes, ordines, genera, species, adiectis synonymis, locis, descriptionibus, observationibus. Kortii, Flensvrgi et Lipsiae [= Flensburg & Leipzig], [32+] 832 pp. <https://doi.org/10.5962/bhl.title.36510>
- Fabricius JC (1805) Systema antiatorum: secundum ordines, genera, species, adiectis synonymis, locis, observationibus, descriptionibus. C. Reichard, Brunsvigae [=Brunswick], 15–372. [1 + 30 pp.] <https://doi.org/10.5962/bhl.title.15806>
- Forsskål P (1775) Descriptiones animalium avium, amphibiorum, piscium, insectorum, vermium; in itinere orientali observavit. Molleri, Havniae [=Copenhagen], [19 +] 34 [+ 164 pp.] <https://doi.org/10.5962/bhl.title.2154>
- Frey R (1946) Übersicht der Gattungen der Syrphiden-Unterfamilie Syrphinae (Syrphine + Bacchinae). Notulae Entomologicae 25[1945]: 152–172.
- van der Goot VS (1964) Fluke's catalogue of Neotropical Syrphidae (Insects, Diptera), a critical study with an appendix on new names in Syrphidae. Beaufortia 10: 212–221.
- Guérin-Méneville FE (1829–1844) Insectes. Iconographie du règne animal de G. Cuvier: ou, Représentation d'après nature de l'une des espèces les plus et souvent non encore figurées de chaque genre d'animaux. JB Bailliere, Paris, 576 pp. [104 pls] <https://doi.org/10.5962/bhl.title.10331>
- Han HY, Choi DS (2001) Family Syrphidae. Economic Insects of Korea 15. Insecta Koreana 22(Supplement): 1–224.
- Hardy DE (1952) Notes and exhibitions. *Ischiodon penicillatus* (Hull). Proceedings of the Hawaiian Entomological Society 14: 363.
- Huang CM, Cheng XY, Yang JK (1996) Diptera: Syrphidae. In: Xue WQ, Chao JM (Eds) Flies of China. Liaoning Science and Technology Press, Shenyang, 118–223.
- Huang C, Cheng X (2012) Diptera: Syrphidae. Fauna Sinica, Insecta vol. 50. Science Press, Neijing, 852 pp. [8 pls]
- Kazerani F, Talebi AA, Gilasian E (2013) An annotated checklist of the subfamily Syrphinae (Diptera: Syrphidae) of Iran. Entomofauna 34: 517–556. [https://www.zobodat.at/pdf/ENT\\_0034\\_0517-0556.pdf](https://www.zobodat.at/pdf/ENT_0034_0517-0556.pdf)
- Kertesz K (1899) Verzeichnis einiger, von L. Biró in Neu-Guinea und am Malayischen Archipel gesammelten Dipteren. Természetrajzi Füzetek kiadja a Magyar nemzeti Muzeum 22: 173–195. <https://biodiversitylibrary.org/page/13453822>
- Khosravian Z, Sadeghi H, Ssymank A (2015) Hover-flies (Diptera: Syrphidae) of Kerman Province, Iran. Far Eastern Entomologist 290: 1–12. <http://www.biosoil.ru/Files/FEE/00000454.pdf>
- Láska P, Pérez-Bañón C, Mazánek L, Rojo S, Ståhls G, Marcos-García MA, Bičík V, Dušek J (2006) Taxonomy of the genera *Scaeva*, *Simosyrphus* and *Ischiodon* (Diptera: Syrphidae): descriptions of immature stages and status of taxa. European Journal of Entomology 103(3): 637–655. <https://doi.org/10.14411/eje.2006.085>
- Le Peletier ALM, Serville JGA (1828) Syrphe. In: Latreille PA, et al. (Eds) Encyclopédie méthodique. Histoire naturelle. Entomologie, ou histoire naturelle des crustacés, des arach-

- nids et des insects. Tome 10. Agasse, Paris, 511–526. <https://biodiversitylibrary.org/page/24404110>
- Linnaeus C (1758) Systema naturae, sive regna tria naturae systematicae proposita per classes, ordines, genera, & species. Ed. 10, Vol. 1. L. Salvii, Holmiae [= Stockholm], 824 pp. <https://doi.org/10.5962/bhl.title.877>
- Loew H (1840) Bemerkungen über die in der Posener Gegend einheimischen Arten mehrerer Zweiflugler-Gattungen. [Zu der] öffentlichen Prüfung der Schuler des Königlichen Friedrich-Wilhelms-Gymnasiums zu Posen 1840, 40 pp. [1 pl]
- Macquart PJM (1842) Syrphids, Syrphidae. In: Dipteres exotiques nouveaux ou peu connus. Tome deuxième. 2e partie. Mémoires de la Société royale des sciences, de l'agriculture et des arts, de Lille 1841(1): 65–200. [22 pls] <https://biodiversitylibrary.org/page/15976657>
- Macquart PJM (1846) Dipteres exotiques nouveaux ou peu connus. Supplement. [1]. Mémoires de la Société royale des sciences, de l'agriculture et des arts, de Lille 1844: 133–364. [20 pls] <https://biodiversitylibrary.org/page/35273814>
- Macquart PJM (1855). Dipteres exotiques nouveaux ou peu connus. 5.e supplement. Mémoires de la Société royale des sciences, de l'agriculture et des arts, de Lille (2) 1: 25–156 [7 pls] <https://biodiversitylibrary.org/page/35274626>
- Matsumura S, Adachi J (1919) Synopsis of the economic Syrphidae of Japan. Pt. III. [sic][=IV]. The Entomological Magazine, Kyoto 3: 128–144. [pl. 3]
- Matsumura S (1916) [Thousand insects of Japan. Additamenta] – Vol. 2 (Diptera). Keisei-sha, Tokyo, 185–474. [16–25 pls; In Japanese] <https://biodiversitylibrary.org/page/34541910>
- Mengual X (2015) *Simosyrphus grandicornis* (flower fly). Invasive Species Compendium. The Centre for Agriculture and Bioscience International (CABI). <http://www.cabi.org/isc/datasheet/50061> [Accessed: 27/03/2018]
- Mengual X, Stähls G, Láska P, Mazánek L, Rojo S (2018) Molecular phylogenetics of the predatory lineage of flower flies *Eupeodes-Scaeva* (Diptera: Syrphidae), with the description of the Neotropical genus *Austroscaeva* gen. nov. Journal of Zoological Systematics and Evolutionary Research 56(2): 148–169. <https://doi.org/10.1111/jzs.12212>
- Naderloo M, Pashaei Rad Sh, Taghaddosi MV (2013) Faunistic study on hover flies (Diptera: Syrphidae) in the eastern part of Zanjan province, Iran. Journal of Entomological Research 4(4): 313–323. [http://jer.iau-arak.ac.ir/article\\_524107\\_83a4c1375d5ec2ee85943b6d4c0fcba5.pdf](http://jer.iau-arak.ac.ir/article_524107_83a4c1375d5ec2ee85943b6d4c0fcba5.pdf)
- Nayar JL (1978) A new species of *Ischiodon* (Syrphidae, Diptera) from Libya. Polskie Pismo Entomologiczne 48: 413–416.
- Rojo S, Gilbert F, Marcos-García MA, Nieto JM, Mier MP (2003) A world review of predatory hoverflies (Diptera, Syrphidae: Syrphinae) and their prey. CIBIO Ediciones, Alicante, 319 pp.
- Sack P (1913) H. Sauter's Formosa-Ausbeute. Syrphiden I. (Dipt.). Entomologische Mitteilungen 2: 1–10. [https://www.zobodat.at/pdf/Entomologische-Mitteilungen\\_2\\_1913\\_0001-0010.pdf](https://www.zobodat.at/pdf/Entomologische-Mitteilungen_2_1913_0001-0010.pdf)
- Sack P (1932) Syrphidae. In: Lindner E (Ed.) Die Fliegen der Palaearktischen Region, IV/6. Schweizerbart, Stuttgart, 451 pp.
- Sarıbüyük S (2014) Check list of Turkish flower flies (Diptera: Syrphidae). Munis Entomology & Zoology 9(1): 570–585. <https://www.munisentzool.org/yayin/vol9/issue1/vol9isue1-4595301.pdf>
- Schiner IR (1860) Vorläufiger Commentar zum dipterologischen Theile der "Fauna Austriaca", mit einer näheren Begründung der in derselben aufgenommenen neuen Dipteren-Gattun-

- gen. II. Wiener Entomologische Monatschrift 4: 208–216. <https://biodiversitylibrary.org/page/31290315>
- Shiraki T (1963) Diptera: Syrphidae. Insects Micronesia 13: 129–187. <http://hbs.bishopmuseum.org/pubs-online/pdf/iom13-5syrph.pdf>
- Shorthouse DP (2010) SimpleMappr, an online tool to produce publication-quality point maps. <http://www.simplemappr.net> [Accessed: 08/02/2018]
- Smit JT, van Harten A, Ketelaar R (2017) Order Diptera, family Syrphidae. The hoverflies of the Arabian Peninsula. Arthropod fauna of the UAE 6: 572–612.
- Speight MCD (2017) Species accounts of European Syrphidae, 2017. Syrph the Net, the database of European Syrphidae (Diptera), Vol. 97. Syrph the Net publications, Dublin, 294 pp.
- Speight MCD, Sarthou J-P (2017) StN keys for the identification of the European species of various genera of Syrphidae 2017. Syrph the Net, the database of European Syrphidae (Diptera), Vol. 99. Syrph the Net publications, Dublin, 139 pp.
- Thompson FC (1999) A key to the genera of the flower flies of the Neotropical Region including the descriptions of genera and species and a glossary of taxonomic terms. Contributions to Entomology, International 3: 319–378. <http://hdl.handle.net/10088/17492>
- Thompson FC (2013) Syrphidae. Systema Dipterorum, version 1.5, 13,354 records. <http://www.diptera.org> [Accessed: 08/02/2018]
- Thomson CG (1869) Diptera. Species nova descriptis. In: Virgin CA (Ed.) Kongliga Svenska Fregatten Eugenies Resa omkring jorden under befäl af C.A. Virgin, Åren 1851–1853. Vetenskapliga Iakttagelser på H.M. Konung Oscar den Förstes befallning utgifna af K. Svenska Vetenskaps Akademien. 2 Zoologi, 1 Insecta. P. A. Norstedt & Söner, Stockholm, 443–614. [https://archive.org/details/cbarchive\\_41647\\_dipteraspesnovadescriptsit1857](https://archive.org/details/cbarchive_41647_dipteraspesnovadescriptsit1857)
- Vockeroth JR (1969) A revision of the genera of the Syrphini (Diptera: Syrphidae). Memoirs of the Entomological Society of Canada 62: 1–176. <https://doi.org/10.4039/entm10162fv>
- Walker F (1852) Diptera. Part III. In: Saunders WW (Ed.) Insecta Saundersiana: or characters of undescribed insects in the collection of William Wilson Saunders, Esq., F.R.S., F.L.S., &c. Vol. 1. Van Voorst, London, 157–252. [5–6 pls] <https://biodiversitylibrary.org/page/42183752>
- Wiedemann CRW (1830) Aussereuropäische Zweiflügelige Insekten, zweiter Theil. Schulz, Hamm, [xii +] 684 pp. [5 pls] <https://doi.org/10.5962/bhl.title.14603>

## Supplementary material I

**Table S1. Localities for the specimens of *Ischiodon astales* Mengual, sp. n. (Geographical coordinates)**

Author: Ximo Mengual

Data type: occurrence

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/AfrInvertebr.59.24461.suppl1>